

Part I.

I - Planning Concepts to Numbers

1. Introduction

There are three kinds of lies: lies, damned lies, and statistics.

Anonymous, but popularized by Mark Twain.

SEATTLE Local News – Two Seconds Cost \$74 Million.¹

That amount was set aside to improve the Mercer Mess. Lanes were added. Signal capacity was improved.

Now GPS navigator TomTom, which tracks drivers using its app, says the average time through the corridor during the peak morning commute 7 minutes, 50 seconds before the Mercer Mess construction.

And now the travel time through Mercer is 7 minutes and 48 seconds. That's right. An improvement of 2 seconds.

¹ \$74 million later, Mercer Mess is 2 seconds faster, KOMO News 2016; case from [callingbullshit](#).

Way of Knowing

Epistemology is a branch of philosophy, which is about the theory of knowledge. For example, they ask how we know what we know and what we can regard as acceptable knowledge.




Figure 1.1: Some ways of knowing: How do we ‘know’ that this tree exists?

Research is a Process of Knowing

Science is a way of knowing while research is the process of finding out the answers, correct or incorrect. Or broadly speaking, research is a process that we gather information systematically to generate new insights and/or answer some problems. For example, we can have:

- Problem: How many homeless individuals are living in King County?
- Research: Count each homeless individual within King County one by one.

This research approach is obviously infeasible. But any other methodologies we use would be an estimation of the real count. Every decision researchers made would influence the final results. The [King County Regional Homelessness Authority's five-year plan \(P.8\)](#) reports five different counts from five different agencies:

Measure	Lead agency	Individuals
2020 Point in Time Count	All Home	11,751
2022 Point in Time Count	KCRHA	13,368
2023 Snapshot Report	WA Department of Commerce	34,866
2020 Cross-Systems Analysis	King County Dept. of Community and Health Services	40,871
2022 Housing Gaps Analysis	WA Dept. of Commerce and Cloudburst Consulting	53,532

Table 1.1: Number of homeless individuals in King County based on different research

The largest is 4.6 times the smallest. We cannot say any of them is a mistake. Each answers a slightly different question. They are counts from different time windows, different definitions of homeless individuals, different data-collection machinery, and different methodologies.

However, as you can imagine, planners will make different decisions, or even opposite decisions if they just use numbers without thinking deeper about some critical questions. Research is not the pursuit of the number; it is the practice of making these choices reasonable and explicit, so that someone else can see what you did, then agree or disagree with it precisely.

Quantitative vs. Qualitative

Data is a collection of facts, numbers, words, observations, or other useful information. We can generally classify data by whether it can be measured/represented using numbers.

Data Types	Methods
Quantitative Data: values that can be measured numerically, such as population, housing prices.	Statistical modeling, simulations, etc.
Qualitative Data: descriptive and non-numerical, capturing characteristics, concepts, or experiences that numbers cannot measure, such as an interview transcript or a survey response.	Case study, text analysis, certain statistical methods, etc.

Table 1.2: Quantitative and qualitative data

Research Paradigms

However, quantitative or qualitative research refers more to the way we analyze data and approach the questions, rather than to the data alone. This taxonomy represents two paradigms of scientific discovery: **positivist** (associated more with

quantitative methods) and **constructionist** (associated more with qualitative methods) (Gliner et al. 2017).

Paradigm	Positivist	Constructionist
The nature of reality	Single reality	Multiple constructed realities
Relationship of knower to known	Researchers are objective	Researchers cannot be totally objective; in fact, researchers and participants interact
Possibility of generalization	Could be generalized	Everything is contextually bound
Possibility of causal linkage	Could at least partially determined	Cannot distinguish cause and effect
The role of value	Research is objective and value free	Bound by choice, values, theory, and conflict

Table 1.3: Positivist and constructionist paradigm

Not all quantitative research is purely positivist, and qualitative research is constructionist. This table reflects the relative differences, and we normally find that most research in planning is a mix of the two. Also, you may realize that quantitative research could use qualitative data (e.g., encoding an interview scripts to find statistical relationship), which we call **mixed method**. This class is focusing on quantitative analysis and URBAN 518 (Autumn quarter 2nd year of MUP) will focus on qualitative analysis.

Math Review

Order of Operations

Evaluate an expression in this order: parentheses, exponents, multiplication and division, then addition and subtraction. Multiplication and division rank equally and are worked left to right, as are addition and subtraction.

$$4^3 \times 3^2 - 10 + 27/3 = 64 \times 9 - 10 + 9 = 575$$

Fractions

A fraction is a **numerator** over a **denominator**. To multiply, multiply numerators and denominators separately; to divide, multiply by the reciprocal:

$$\frac{1}{2} \times \frac{3}{4} = \frac{1 \times 3}{2 \times 4} = \frac{3}{8}, \quad \frac{1}{2} \div \frac{3}{4} = \frac{1}{2} \times \frac{4}{3} = \frac{2}{3}$$

Adding or subtracting requires a common denominator: $\frac{1}{3} + \frac{1}{4} = \frac{4}{12} + \frac{3}{12} = \frac{7}{12}$.

Every rate is a fraction, and every unit conversion is a fraction multiplication: one-third of an acre per 100 people is $\frac{1/3}{100} \times 1000 = 3.33$ acres per 1,000 people.

Solving for an Unknown

Apply the same operation to both sides of an equation until the unknown stands alone. Both sides stay equal because each received the same change. A standard of

8 acres per 1,000 residents, at a projected population of 806,800, requires a acres:

$$\frac{a}{806,800} \times 1000 = 8 \implies a \times 1000 = 8 \times 806,800 \implies a = \frac{8 \times 806,800}{1000} = 6,454.4$$

Lines and Slope

A line is written $y = mx + b$. The **slope** m is the change in y per one-unit change in x ; the **intercept** b is the value of y when $x = 0$. Given two points (x_1, y_1) and (x_2, y_2) :

$$m = \frac{y_2 - y_1}{x_2 - x_1}, \quad b = y_1 - mx_1$$

The slope is the rise over the run, and its sign says whether y increases or decreases as x increases. Seattle's population was 608,660 in 2010 and 686,800 in 2016, so the slope through those two points is $(686,800 - 608,660)/(2016 - 2010) \approx 13,023$ people per year: the average annual increment implied by a straight line.




Figure 1.2: Anatomy of a line. The slope is the rise over the run; the intercept is the height of the line at $x = 0$.

Percent versus Percentage Point

A **percentage point** is an arithmetic difference between two percentages. A **percent change** is a relative change. Moving from 20% to 25% is a rise of 5 percentage points and a rise of 25 percent.

The Seattle Parks and Open Space Plan contains the sentence that:

| about 23% of Seattleites are aged 25 to 34, “which is up 21% from 2010”.

This is a relative increase in the count, not a shift of 21 percentage points in the share. Read the wrong way, that sentence implies a 2010 share of 2%.

Growth Rates

Given a starting value P_0 and an ending value P_1 separated by n years, the **compound annual growth rate (CAGR)** is the constant yearly rate that would carry one to the other:

$$\text{CAGR} = \sqrt[n]{\frac{P_1}{P_0}} - 1 = \left(\frac{P_1}{P_0}\right)^{1/n} - 1$$

If projecting, a **linear** assumption adds a fixed number of people each year while a **geometric** assumption multiplies by a fixed factor. Over a twenty-year horizon, the two diverge substantially.




Figure 1.3: Linear and geometric growth through Seattle's 2010 and 2016 populations (608,660 and 686,800). The linear path adds 13,023 people a year; the geometric path grows 2.03% a year, the CAGR.

Weighted Averages

The mean of a set of subgroup means is not the mean of the whole unless the subgroups are the same size. With groups of size n_j and group means $\bar{y}_j$, the correct combination is

$$\bar{y} = \frac{n_1}{n} \bar{y}_1 + \frac{n_2}{n} \bar{y}_2 + \cdots + \frac{n_k}{n} \bar{y}_k, \quad n = n_1 + n_2 + \cdots + n_k$$

Example 1.1. Calculate the average age of two groups.




Figure 1.4: Two groups of 30 and 10 people and their average ages. Each group's area is its share of the whole.

The overall average age needs to be weighted by the proportion of people, not simply the average of the two group averages.

$$\text{Overall Age} = \frac{30}{30 + 10} \times 20 + \frac{10}{30 + 10} \times 40 = 25$$

Notation and Calculus OPTIONAL

Function OPTIONAL

A function $f(x)$ assigns one output to each input. The domain is the set of allowed inputs; the range is the set of possible outputs.

For example, $f(x) = 2x + 1$ maps the input $x = 3$ to the output $f(3) = 2 \times 3 + 1 = 7$. Its domain and range are all real numbers. The function $g(x) = 1/x$ is undefined at $x = 0$, so its domain is every $x \neq 0$.

Summation OPTIONAL

$\sum_{i=1}^n x_i = x_1 + x_2 + \cdots + x_n$. The index i runs over the cases; n is the number of cases.

Note that $\sum x_i^2$ and $(\sum x_i)^2$ are different: for the values 1, 2, 3, $\sum x_i^2 = 14$ while $(\sum x_i)^2 = 36$.

Factorial OPTIONAL

$n! = n(n-1)(n-2) \cdots 1$, with $0! = 1$. Example: $4! = 4 \times 3 \times 2 \times 1 = 24$. Appears in counting formulas for arrangements and combinations.

Why $0! = 1$.

Based on the definition, we can know the factorial in terms of the next one:

$$n! = \frac{(n+1)!}{n+1}$$

So that

$$0! = \frac{(0+1)!}{0+1} = \frac{1!}{1} = \frac{1}{1} = 1$$

Defining **$0! = 1$** is the only way to keep this mathematical logic consistent.

Exponents and Roots OPTIONAL

a^n means a multiplied by itself n times; a is the **base** and n the **exponent**. Exponents need not be whole numbers.

$$a^m a^n = a^{m+n}, \quad (a^m)^n = a^{mn}, \quad \frac{a^m}{a^n} = a^{m-n}, \quad a^{-n} = \frac{1}{a^n}, \quad a^0 = 1$$

A **root** is a fractional exponent: $a^{1/2} = \sqrt{a}$, and $a^{1/n}$ is the n th root of a .

Logarithm OPTIONAL

The inverse of exponentiation: if $y = a^x$, then $\log_a y = x$.

Some rules: $\log(xy) = \log x + \log y$; $\log(x/y) = \log x - \log y$; $\log(x^k) = k \log x$.

In these notes $\log$ denotes the natural logarithm, base $e \approx 2.718$, with inverse $\exp(x) = e^x$.

Reference: [What's so special about Euler's number e?](#) 3Blue1Brown

Limit OPTIONAL

$\lim_{x \rightarrow a} f(x) = L$: the value of $f(x)$ can be made close to L by taking x close enough to a , with $x \neq a$. The target a is a number or $\pm\infty$, and “close enough” reads differently in each case:

- $x \rightarrow \infty$: x large enough. Written $\lim_{n \rightarrow \infty}$, it underlies “as n grows” statements.
- $x \rightarrow -\infty$: x negative enough.
- $x \rightarrow a$ for a number a : $|x - a|$ small enough. The derivative below takes $h \rightarrow 0$, that is, $|h|$ small enough.

For $f(x) = 1/x$, $\lim_{x \rightarrow \infty} 1/x = 0$ and $\lim_{x \rightarrow -\infty} 1/x = 0$ (Figure 1.5). The limit is the value approached, not a value reached.




Figure 1.5: The limit of $1/x$ as $x \rightarrow \infty$ or $x \rightarrow -\infty$ is 0.

Matrix OPTIONAL

An $n \times k$ matrix is a rectangular array with n rows and k columns; a vector is a single column. A data frame is a matrix with named columns.

$$\mathbf{X} = \begin{bmatrix} x_{11} & x_{12} & \cdots & x_{1k} \\ x_{21} & x_{22} & \cdots & x_{2k} \\ \vdots & \vdots & \ddots & \vdots \\ x_{n1} & x_{n2} & \cdots & x_{nk} \end{bmatrix}, \quad \mathbf{y} = \begin{bmatrix} y_1 \\ y_2 \\ \vdots \\ y_n \end{bmatrix}$$

The entry x_{ij} sits in row i and column j . In a data set, row i is case i and column j is variable j . The vector $\mathbf{y}$ is $n \times 1$: one value per case.

Derivative OPTIONAL

$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$: the changing rate of f at x , the slope of the tangent line at that point. Common rules:

- $\frac{d}{dx} x^n = nx^{n-1}$
- $\frac{d}{dx} e^x = e^x$
- $\frac{d}{dx} \ln x = 1/x$
- $[f(x)g(x)]' = f'(x)g(x) + f(x)g'(x)$ (product rule)
- $[f(g(x))]' = f'(g(x))g'(x)$ (chain rule)

For $f(x) = x^2$ at $x = 1$, the line through $(1, 1)$ and $(1 + h, (1 + h)^2)$ has slope

$$\frac{(1 + h)^2 - 1^2}{h} = \frac{2h + h^2}{h} = 2 + h.$$

As $h \rightarrow 0$ the slope goes to 2, which matches: $f'(1) = 2 \times 1 = 2$ (Figure 1.6).




Figure 1.6: The derivative of $f(x) = x^2$ at $x = 1$. Left: lines through $(1, 1)$ and $(1 + h, (1 + h)^2)$ have slope $2 + h$.

Integral OPTIONAL

$\int_a^b f(x) dx$ is the area under f between a and b .

An **antiderivative** of f is any F with $F'(x) = f(x)$, written $\int f(x) dx = F(x) + C$; so that $\int_a^b f(x) dx = F(b) - F(a)$. Some basic forms:

- $\int x^n dx = \frac{x^{n+1}}{n+1} + C$ for $n \neq -1$
- $\int \frac{1}{x} dx = \ln|x| + C$
- $\int e^x dx = e^x + C$

For the constant function $f(x) = 5$, the antiderivative $F(x) = 5x + C$ gives: $F(b) - F(a) = 5b - 5a$. In another way, the region under f between a and b is a rectangle of height 5 and width $b - a$ (Figure 1.7):




Figure 1.7: The integral of the constant function $f(x) = 5$ from a to b is the shaded rectangle, with area $5(b - a)$.

Example 1.2. Density and distribution function. For a variable X , we can define F as the accumulated area below the curve from $-\infty$ to x_0 (an integral operation):

$$F(x_0) = \int_{-\infty}^{x_0} f(t) dt = P(X \leq x_0)$$

Differentiating undoes the accumulation and returns the $f(x_0)$, which is a rate how fast F is accruing at x_0 :

$$f(x_0) = F'(x_0) = \lim_{h \rightarrow 0} \frac{F(x_0 + h) - F(x_0)}{h}.$$




Figure 1.8: Differentiation and integration are one relationship read in two directions. Integrating the density f up to x_0 gives the area $F(x_0) = P(X \leq x_0)$; differentiating F at x_0 gives back the height $f(x_0)$, which is the slope.